

Notes in vaccination (Please read before inoculation)

Updated: October 26, 2016

Please do to inoculate when good physical condition ●.

- BCG · MR (measles-rubella), varicella, such as those that are not older than 27 days in response to the vaccination of live vaccines can not be inoculated.
- can not be inoculated person who has not passed more than six days in response to the vaccination of inactivated vaccine, such as four types mixed and Japanese encephalitis.
- vaccination day to avoid the extreme exercise, please observe the rest.
- Please have is maintaining the cleanliness was inoculated.
- After inoculation, Dari damaged or inoculation site becomes red, but you may mild fever, will be progressively better during the normal two to three days. However, when there is a high fever and vomiting spasms other abnormal symptoms, please consult a doctor as soon as possible.

To receive the vaccination in Koganei, you will need vaccination coupon of each vaccination. By the time the child is age 2 months, will be sent as a "vaccination notebook" from the city. Procedures, etc. is not required.

You were moving in, If you do not have a vaccination coupon of Koganei City, whether you send it to "vaccination completed notification form" (postcard) the health department, we ask that you contact. We will send a vaccination coupon.


<Medical institutions>

Since the vaccination carried out medical institution depends on the vaccination, please refer to the "city vaccination implementation list" below.

In addition to the Koganei, even in Mitaka City, Musashino City, Fuchu, Tachikawa, Akishima City, Kodaira City, Higashimurayama-Kokubunji National City, Komae City, Higashiyamato City, Kiyose City, Musashimurayama, There is a medical institution to receive the vaccination.

Other city, when it is inoculated with outside Tokyo, you will need a separate procedure. For more information, please contact the health department.

 [City pediatric immunization practice medical institution list \(PDF: 172KB\)](#)

 [Childhood routine immunization implementation list \(PDF: 175KB\)](#)

<Health Relief System>

Vaccination, but is useful in order to prevent the infection, on the other hand there is extremely rare, is seen the occurrence of health damage. If the health damage caused by regular vaccination occurs, there is a health damage relief system after vaccination with the country.

The side reaction, fever and that occurs after inoculated with the vaccine, relatively well lighter side reaction and seen, such as swelling of the inoculation site, there is a serious side reaction to an extremely rare occurrence, such as encephalitis and neuropathy.

However, the side reactions, the vaccine, such as accidental infection that developed around the same time may not be the cause. In the vaccination after the health damage relief system, whether or not due to vaccination reviewed individually, it will be paid in the case that has been certified to be due to vaccination.

 [Ministry of Health, Labour and Welfare website \(for health Relief System\) \(external site\)](#)